



Maine Medical Association

Robert J. Schlager, MD, President | Amy Madden, MD, President-Elect | Karen Saylor, MD, Chair, Board of Directors
Andrew B. MacLean, JD, Interim CEO | Peter P. Michaud, JD, RN, General Counsel

STATEMENT ON PHYSICIAN-ASSISTED SUICIDE

APRIL 2019

Since legislation was introduced before the 128th Maine Legislature (2017-2018), the Maine Medical Association (MMA) has been re-evaluating its position in opposition to physician-assisted suicide (also referred to as physician-assisted death) established by a 2001 resolution of its membership.

The MMA Board of Directors formed an ad hoc committee of members interested in the issue to review its position and to make a recommendation to the Board. The ad hoc committee directed the MMA to solicit the opinions of members through several polls and considered the positions of other state medical societies and the Council on Ethical & Judicial Affairs (CEJA) of the American Medical Association (AMA). The polling results showed that physicians are divided on the issue.

Upon careful deliberation, the ad hoc committee noted that the AMA Code of Medical Ethics already provides guidance on this issue and recommended to the Board that the MMA provide information and seek to educate members, elected officials and policymakers but take no position in support of or opposition to physician-assisted death in any proposed legislation before the Maine legislature or proposed referenda question before Maine voters. Whether to establish in Maine law, a pathway for physician-assisted death is a policymaking decision for Maine voters and their elected representatives to make.

While MMA will take no position on the central issue of physician-assisted death in such proposals, it will testify as needed on any provisions of such initiatives that might adversely affect patients or physicians and it will continue its long-standing advocacy for the best possible end-of-life care, including hospice care, for all residents of both urban and rural parts of our state. MMA urges its members to exercise their individual voting rights in accordance with their personal conscience, values, and interpretation of two key provisions of the AMA Code of Medical Ethics and the most recent policy statement of the AMA's CEJA described further below.

Upon vote of the Board of Directors during its regular meeting on Wednesday, April 24, 2019, the MMA offers the following three documents (attached) as the most current, relevant guidance for members as they refine their personal views on this challenging societal issue.

1. Code of Medical Ethics Opinion 5.7, *Physician-Assisted Suicide*;
2. Code of Medical Ethics Opinion 1.1.7, *Physician Exercise of Conscience*; and
3. CEJA Report 2-1-18, the most recent AMA policy statement on the issue.

The MMA Board has concluded that the AMA CEJA is not likely to amend or eliminate the two Code of Medical Ethics statements cited above and that CEJA Report 2-1-18 acknowledges, but does not attempt to resolve, the ethical and philosophical tensions inherent in this societal debate. The MMA will continue to promote discussion of the issue within the profession.

Code of Medical Ethics

5.7 Physician-Assisted Suicide

Topic: Code of Medical Ethics **Policy Subtopic:** Opinions on Caring for Patients at the End of Life

Meeting Type: NA

Year Last Modified: 2017

Action: NA

Type: Code of Medical Ethics

Council & Committees: NA



Physician-assisted suicide occurs when a physician facilitates a patient's death by providing the necessary means and/or information to enable the patient to perform the life-ending act (e.g., the physician provides sleeping pills and information about the lethal dose, while aware that the patient may commit suicide).

It is understandable, though tragic, that some patients in extreme duress—such as those suffering from a terminal, painful, debilitating illness—may come to decide that death is preferable to life. However, permitting physicians to engage in assisted suicide would ultimately cause more harm than good.

Physician-assisted suicide is fundamentally incompatible with the physician's role as healer, would be difficult or impossible to control, and would pose serious societal risks.

Instead of engaging in assisted suicide, physicians must aggressively respond to the needs of patients at the end of life. Physicians:

- (a) Should not abandon a patient once it is determined that cure is impossible.
- (b) Must respect patient autonomy.
- (c) Must provide good communication and emotional support.
- (d) Must provide appropriate comfort care and adequate pain control.

AMA Principles of Medical Ethics: I,IV

The Opinions in this chapter are offered as ethics guidance for physicians and are not intended to establish standards of clinical practice or rules of law.

Policy Timeline

Issued: 2016

Code of Medical Ethics

1.1.7 Physician Exercise of Conscience

Topic: Code of Medical Ethics **Policy Subtopic:** Opinions on Patient-Physician Relationships (1.1 Responsibilities of Physicians & Patients)
Meeting Type: NA **Year Last Modified:** 2017
Action: NA **Type:** Code of Medical Ethics
Council & Committees: NA



Physicians are expected to uphold the ethical norms of their profession, including fidelity to patients and respect for patient self-determination. Yet physicians are not defined solely by their profession. They are moral agents in their own right and, like their patients, are informed by and committed to diverse cultural, religious, and philosophical traditions and beliefs. For some physicians, their professional calling is imbued with their foundational beliefs as persons, and at times the expectation that physicians will put patients' needs and preferences first may be in tension with the need to sustain moral integrity and continuity across both personal and professional life.

Preserving opportunity for physicians to act (or to refrain from acting) in accordance with the dictates of conscience in their professional practice is important for preserving the integrity of the medical profession as well as the integrity of the individual physician, on which patients and the public rely. Thus physicians should have considerable latitude to practice in accord with well-considered, deeply held beliefs that are central to their self-identities.

Physicians' freedom to act according to conscience is not unlimited, however. Physicians are expected to provide care in emergencies, honor patients' informed decisions to refuse life-sustaining treatment, and respect basic civil liberties and not discriminate against individuals in deciding whether to enter into a professional relationship with a new patient.

In other circumstances, physicians may be able to act (or refrain from acting) in accordance with the dictates of their conscience without violating their professional obligations. Several factors impinge on the decision to act according to conscience. Physicians have stronger obligations to patients with whom they have a patient-physician relationship, especially one of long standing; when there is imminent risk of foreseeable harm to the patient or delay in access to treatment would significantly adversely affect the patient's physical or emotional well-being; and when the patient is not reasonably able to access needed treatment from another qualified physician.

In following conscience, physicians should:

- (a) Thoughtfully consider whether and how significantly an action (or declining to act) will undermine the physician's personal integrity, create emotional or moral distress for the physician, or compromise the physician's ability to provide care for the individual and other patients.

- (b) Before entering into a patient-physician relationship, make clear any specific interventions or services the physician cannot in good conscience provide because they are contrary to the physician's deeply held personal beliefs, focusing on interventions or services a patient might otherwise reasonably expect the practice to offer.
- (c) Take care that their actions do not discriminate against or unduly burden individual patients or populations of patients and do not adversely affect patient or public trust.
- (d) Be mindful of the burden their actions may place on fellow professionals.
- (e) Uphold standards of informed consent and inform the patient about all relevant options for treatment, including options to which the physician morally objects.
- (f) In general, physicians should refer a patient to another physician or institution to provide treatment the physician declines to offer. When a deeply held, well-considered personal belief leads a physician also to decline to refer, the physician should offer impartial guidance to patients about how to inform themselves regarding access to desired services.
- (g) Continue to provide other ongoing care for the patient or formally terminate the patient-physician relationship in keeping with ethics guidance.

AMA Principles of Medical Ethics: I,II,IV,VI,VIII,IX

The Opinions in this chapter are offered as ethics guidance for physicians and are not intended to establish standards of clinical practice or rules of law.

Policy Timeline

Issued: 2016

REPORT OF THE COUNCIL ON ETHICAL AND JUDICIAL AFFAIRS*

CEJA Report 2-I-18

Subject: Study Aid-in-Dying as End-of-Life Option
(Resolution 15-A-16)
The Need to Distinguish “Physician-Assisted Suicide” and “Aid in Dying”
(Resolution 14-A-17)

Presented by: James E. Sabin, MD, Chair

Referred to: Reference Committee on Amendments to Constitution and Bylaws
(Todd M. Hertzberg, MD, Chair)

1 At the 2016 Annual Meeting, the House of Delegates referred Resolution 15-A-16, “Study Aid-in-
2 Dying as End-of-Life Option,” presented by the Oregon Delegation, which asked:
3

4 That our American Medical Association (AMA) and its Council on Judicial and Ethical
5 Affairs (CEJA), study the issue of medical aid-in-dying with consideration of (1) data
6 collected from the states that currently authorize aid-in-dying, and (2) input from some of
7 the physicians who have provided medical aid-in-dying to qualified patients, and report
8 back to the HOD at the 2017 Annual Meeting with recommendation regarding the AMA
9 taking a neutral stance on physician “aid-in-dying.”
10

11 At the following Annual Meeting in June 2017, the House of Delegates similarly referred
12 Resolution 14-A-17, “The Need to Distinguish between ‘Physician-Assisted Suicide’ and ‘Aid in
13 Dying’” (presented by M. Zuhdi Jasser, MD), which asked that our AMA:
14

15 (1) as a matter of organizational policy, when referring to what it currently defines as
16 ‘*Physician Assisted Suicide*’ avoid any replacement with the phrase ‘*Aid in Dying*’ when
17 describing what has long been understood by the AMA to specifically be ‘*Physician Assisted*
18 *Suicide*’; (2) develop definitions and a clear distinction between what is meant when the AMA
19 uses the phrase ‘*Physician Assisted Suicide*’ and the phrase ‘*Aid in Dying*’; and (3) fully utilize
20 these definitions and distinctions in organizational policy, discussions, and position statements
21 regarding both ‘*Physician Assisted Suicide*’ and ‘*Aid in Dying*.’
22

23 This report by the Council on Ethical and Judicial Affairs addresses the concerns expressed in
24 Resolutions 15-A-16 and 14-A-17. In carrying out its review of issues in this area, CEJA reviewed
25 the philosophical and empirical literature, sought input from the House of Delegates through an I-
26 16 educational program on physician-assisted suicide, an informal “open house” at A-17, and its I-
27 17 Open Forum. The council wishes to express its sincere appreciation for participants’
28 contributions during these sessions and for additional written communications received from
29 multiple stakeholders, which have enhanced its deliberations.

* Reports of the Council on Ethical and Judicial Affairs are assigned to the Reference Committee on Amendments to Constitution and Bylaws. They may be adopted, not adopted, or referred. A report may not be amended, except to clarify the meaning of the report and only with the concurrence of the Council.

1 The council observes that the ethical arguments advanced today supporting and opposing
2 "physician-assisted suicide" or "aid in dying" are fundamentally unchanged from those examined
3 in CEJA's 1991 report on this topic [1]. The present report does not rehearse these arguments again
4 as such. Rather, it considers the implications of the legalization of assisted suicide in the United
5 States since the adoption of Opinion E-5.7, "Physician-Assisted Suicide," in 1994.

6 7 "ASSISTED SUICIDE," "AID IN DYING," OR "DEATH WITH DIGNITY"?

8
9 Not surprisingly, the terms stakeholders use to refer the practice of physicians prescribing lethal
10 medication to be self-administered by patients in many ways reflect the different ethical
11 perspectives that inform ongoing societal debate. Proponents of physician participation often use
12 language that casts the practice in a positive light. "Death with dignity" foregrounds patients'
13 values and goals, while "aid in dying" invokes physicians' commitment to succor and support.
14 Such connotations are visible in the titles of relevant legislation in states that have legalized the
15 practice: "Death with Dignity" (Oregon, Washington, District of Columbia), "Patient Choice and
16 Control at the End of Life" (Vermont), "End of Life Options" (California, Colorado), "Our Care
17 Our Choice Act" (Hawaii), and in Canada's "Medical Aid in Dying."

18
19 Correspondingly, those who oppose physician provision of lethal medications refer to the practice
20 as "physician-assisted suicide," with its negative connotations regarding patients' psychological
21 state and its suggestion that physicians are complicit in something that, in other contexts, they
22 would seek to prevent. The language of dignity and aid, critics contend, are euphemisms [2]; their
23 use obscures or sanitizes the activity. In their view such language characterizes physicians' role in
24 a way that risks construing an act that is ethically unacceptable as good medical practice [3]. Still
25 others, meanwhile, argue that the choice by terminally ill patients to take action to end their own
26 lives with the assistance of their physician is distinct from what is traditionally understood as
27 "suicide" [4].

28
29 The council recognizes that choosing one term of art over others can carry multiple, and not always
30 intended messages. However, in the absence of a perfect option, CEJA believes ethical deliberation
31 and debate is best served by using plainly descriptive language. In the council's view, despite its
32 negative connotations [5], the term "physician assisted suicide" describes the practice with the
33 greatest precision. Most importantly, it clearly distinguishes the practice from euthanasia [1]. The
34 terms "aid in dying" or "death with dignity" could be used to describe either euthanasia or
35 palliative/hospice care at the end of life and this degree of ambiguity is unacceptable for providing
36 ethical guidance.

37 38 COMMON GROUND

39
40 Beneath the seemingly incommensurate perspectives that feature prominently in public and
41 professional debate about writing a prescription to provide patients with the means to end life if
42 they so choose, CEJA perceives a deeply and broadly shared vision of what matters at the end of
43 life. A vision that is characterized by hope for a death that preserves dignity, a sense of the
44 sacredness of ministering to a patient at the end of life, recognition of the relief of suffering as the
45 deepest aim of medicine, and fully voluntary participation on the part of both patient and physician
46 in decisions about how to approach the end of life.

47
48 Differences lie in the forms these deep commitments take in concrete decisions and actions. CEJA
49 believes that thoughtful, morally admirable individuals hold diverging, yet equally deeply held, and
50 well-considered perspectives about physician-assisted suicide that govern how these shared
51 commitments are ultimately expressed. For one patient, dying "with dignity" may mean accepting

1 the end of life however it comes as gracefully as one can; for another, it may mean being able to
2 exercise some measure of control over the circumstances in which death occurs. For some
3 physicians, the sacredness of ministering to a terminally ill or dying patient and the duty not to
4 abandon the patient preclude the possibility of supporting patients in hastening their death. For
5 others, not to provide a prescription for lethal medication in response to a patient's sincere request
6 violates that same commitment and duty. Both groups of physicians base their view of ethical
7 practice on the guidance of Principle I of the *AMA Principles of Medical Ethics*: "A physician
8 shall be dedicated to providing competent medical care, with compassion and respect for human
9 dignity and rights."

10
11 So too, how physicians understand and act on the goals of relieving suffering, respecting
12 autonomy, and maintaining dignity at the end of life is directed by identity-conferring beliefs and
13 values that may not be commensurate. Where one physician understands providing the means to
14 hasten death to be an abrogation of the physician's fundamental role as healer that forecloses any
15 possibility of offering care that respects dignity, another in equally good faith understands
16 supporting a patient's request for aid in hastening a foreseen death to be an expression of care and
17 compassion.

18 19 IRREDUCIBLE DIFFERENCES IN MORAL PERSPECTIVES ON PHYSICIAN-ASSISTED 20 SUICIDE

21
22 How to respond when coherent, consistent, and deeply held beliefs yield irreducibly different
23 judgments about what is an ethically permissible course of action is profoundly challenging. With
24 respect to physician-assisted suicide, some professional organizations—for example, the American
25 Academy of Hospice and Palliative Medicine [6]—have adopted a position of "studied neutrality."
26 Positions of studied neutrality neither endorse nor oppose the contested practice, but instead are
27 intended to respect that there are irreducible differences among the deeply held beliefs and values
28 that inform public and professional perspectives [6,7], and to leave space open for ongoing
29 discussion. Nonetheless, as a policy position, studied neutrality has been criticized as neither
30 neutral or appropriate for organized medicine [8], and as being open to unintended consequences,
31 including stifling the very debate it purports to encourage or being read as little more than
32 acquiescence with the contested practice [9].

33
34 CEJA approaches the condition of irreducible difference from a different direction. In its 2014
35 report on exercise of conscience, the Council noted that "health care professionals may hold very
36 different core beliefs and thus reach very different decisions based on those core beliefs, yet
37 equally act according to the dictates of conscience. For example, a physician who chooses to
38 provide abortions on the basis of a deeply held belief in protecting women's autonomy makes the
39 same kind of moral claim to conscience as does a physician who refuses to provide abortion on the
40 basis of respect for the sanctity of life of the fetus" [10].

41
42 Importantly, decisions taken in conscience are not simply idiosyncratic; they do not rest on
43 intuition or emotion. Rather, such decisions are based on "substantive, coherent, and reasonably
44 stable" values and principles [10]. Physicians must be able to articulate how those values and
45 principles justify the action in question.

46
47 The ethical arguments offered for more than two decades by those who support and those who
48 oppose physician participation in assisted suicide reflect the diverging "substantive, coherent, and
49 reasonably stable" values and principles within the profession and the wider moral community.
50 While supporters and opponents of physician-assisted suicide share a common commitment to
51 "compassion and respect for human dignity and rights" (*AMA Principles of Medical Ethics*, I),

1 they draw different moral conclusions from the underlying principle they share. As psychiatrist
2 Harvey Chochinov observed with respect to the stakeholders interviewed by Canadian Supreme
3 Court's advisory panel on physician-assisted death, "neither those who are strongly supportive nor
4 those who are opposed hold a monopoly on integrity and a genuine concern for the well-being of
5 people contemplating end of life. Equally true: neither side is immune from impulses shaped more
6 by ideology than a deep and nuanced understanding of how to best honor and address the needs of
7 people who are suffering" [11].
8

9 THE RISK OF UNINTENDED CONSEQUENCES

10
11 From the earliest days of the debate, a prominent argument raised against permitting physician-
12 assisted suicide has been that doing so will have adverse consequences for individual patients, the
13 medical profession, and society at large. Scholars have cited the prospect that boundaries will be
14 eroded and practice will be extended beyond competent, terminally ill adult patients; to patients
15 with psychiatric disorders, children; or that criteria will be broadened beyond physical suffering to
16 encompass existential suffering; or that stigmatized or socioeconomically disadvantaged patients
17 will be coerced or encouraged to end their lives. Concerns have also been expressed that permitting
18 the practice will compromise the integrity of the profession, undermine trust, and harm the
19 physicians and other health care professionals who participate; and that forces outside medicine
20 will unduly influence decisions.
21

22 The question whether safeguards—which in the U.S. jurisdictions that permit assisted suicide,
23 restrict the practice to terminally ill adult patients who have decision-making capacity and who
24 voluntarily request assisted suicide, along with procedural and reporting requirements—can
25 actually protect patients and sustain the integrity of medicine remains deeply contested. Some
26 studies have "found no evidence to justify the grave and important concern often expressed about
27 the potential for abuse—namely, the fear that legalized physician-assisted dying will target the
28 vulnerable or pose the greatest risk to people in vulnerable groups" [12], others question whether
29 the available data can in fact support any such conclusions, finding the evidence cited variously
30 flawed [13], inadequate [14], or distorted [15].
31

32 Although cross-cultural comparisons are problematic [16], current evidence from Europe does tell
33 a cautionary tale. Recent findings from studies in Belgium and the Netherlands, both countries that
34 permit euthanasia as well as physician-assisted suicide, mitigate some fears but underscore others
35 [17]. For example, research in the Netherlands has found that "requests characterized by
36 psychological as opposed to physical suffering were more likely to be rejected, as were requests by
37 individuals who lived alone," mitigating fears that "solitary, depressed individuals with potentially
38 reversible conditions might successfully end their lives." At the same time, however, among
39 patients who obtained euthanasia or assisted suicide, nearly 4 percent "reported only psychological
40 suffering." At the level of anecdote, a description of a case of euthanasia in Belgium elicited
41 widespread concern about the emergence of a "slippery slope" [18].
42

43 Studies have also raised questions about how effective retrospective review of decisions to provide
44 euthanasia/assisted suicide is in policing practice [19,20]. A qualitative analysis of cases that Dutch
45 regional euthanasia committees determined had not met legal "due care criteria" found that such
46 reviews focus on procedural considerations and do not "directly assess the actual eligibility" of the
47 patients who obtained euthanasia [19]. A separate study of cases in which psychiatric patients
48 obtained euthanasia found that physicians' reports "stated that psychosis or depression did or did
49 not affect capacity but provided little explanation regarding their judgments" and that review
50 committees "generally accepted the judgment of the physician performing EAS [euthanasia or
51 physician-assisted suicide]" [20]. It remains an open question whether reviews that are not able to

1 assess physicians' reasoning truly offer the protection they are intended to provide. To the extent
2 that reporting and data collection in states that permit physician-assisted suicide have similar
3 limitations, oversight of practice may not be adequate.
4

5 Medicine must learn from this experience. Where physician-assisted suicide is legalized,
6 safeguards can and should be improved—e.g., “[t]o increase safeguards, states could consider
7 introducing multidisciplinary panels to support patients through the entire process, including
8 verifying consent and capacity, ensuring appropriate psychosocial counseling, and discussing all
9 palliative and end-of-life options” [21]. Both the state and the medical profession have a
10 responsibility to monitor ongoing practice in a meaningful way and to address promptly
11 compromises in safeguards should any be discovered. It is equally important that strong practices
12 be identified and encouraged across all jurisdictions that permit physicians to assist suicide. Health
13 care organizations in California and Canada, for example, have shared richly descriptive reports of
14 practices adopted in response to the recent legalization of “aid in dying” in those jurisdictions that
15 seek to address concerns about quality of practice and data collection [22,23].
16

17 Medicine must also acknowledge, however, that evidence (no matter how robust) that there have
18 not yet been adverse consequences cannot guarantee that such consequences would not occur in the
19 future. As a recent commentary noted, “[p]art of the problem with the slippery slope is you never
20 know when you are on it” [17].
21

22 SAFEGUARDING DECISIONS AT THE END OF LIFE 23

24 CEJA has found that just as there are shared commitments behind deep differences regarding
25 physician-assisted suicide, there are also shared concerns about how to understand the available
26 evidence. For example, in the council's recent Open Forum, both proponents and opponents of
27 physician-assisted suicide observed that in the U.S., debate occurs against the backdrop of a health
28 care system in which patients have uneven access to care, including access to high quality end-of-
29 life care. They also noted that patients and physicians too often still do not have the conversations
30 they should about death and dying, and that too few patients are aware of the range of options for
31 end-of-life care, raising concern that many patients may be led to request assisted suicide because
32 they don't understand the degree of relief of suffering state-of-the-art palliative care can offer.
33 Participants who in other respects held very different views concurred as well that patients may be
34 vulnerable to coercion, particularly patients who are in other ways disadvantaged; and expressed
35 concern in common that forces external to medicine could adversely influence practice.
36

37 These are much the same concerns the Institute of Medicine identified in its 2015 report, *Dying in*
38 *America* [24]. They are concerns echoed in a February 2018 workshop on physician-assisted death
39 convened by the National Academies of Science, Engineering and Medicine [25]. They underscore
40 how important it is to understand *why* a patient requests assisted suicide as a starting point for care
41 [26].
42

43 Patient requests for assisted suicide invite physicians to have the kind of difficult conversations that
44 are too often avoided. They open opportunities to explore the patient's goals and concerns, to learn
45 what about the situation the individual finds intolerable and to respond creatively to the patient's
46 needs other than providing the means to end life—by such means as better managing symptoms,
47 arranging for psychosocial or spiritual support, treating depression, and helping the patient to
48 understand more clearly how the future is likely to unfold [5,27]. Medicine as a profession must
49 ensure that physicians are skillful in engaging in these difficult conversations and knowledgeable
50 about the options available to terminally ill patients [28]. The profession also has a responsibility to
51 advocate for adequate resources for end-of-life care [16,28], particularly for patients from

1 disadvantaged groups. The availability of assisted suicide where it is legal must not be allowed to
2 interfere with excellent care at the end of life.

3
4 CONCLUSION

5
6 At the core of public and professional debate, the council believes, is the aspiration that every
7 patient come to the end of life as free as possible from suffering that does not serve the patient's
8 deepest self-defining beliefs and in the presence of trusted companions, including where feasible
9 and when the patient desires, the presence of a trusted physician. As Timothy Quill noted more
10 than 20 years ago, "dying patients do not have the luxury of choosing not to undertake the journey,
11 or of separating their person from their disease" [27]. Decisions about how to approach the end of
12 life are among the most intimate that patients, families, and their physicians make. Respecting the
13 intimacy and the authenticity of those relationships is essential if our common ideal is to be
14 achieved.

15
16 While supporters and opponents of physician-assisted suicide share a common commitment to
17 "compassion and respect for human dignity and rights" (AMA Principles of Medical Ethics, I),
18 they draw different moral conclusions from the underlying principle they share. Where one
19 physician understands providing the means to hasten death to be an abrogation of the physician's
20 fundamental role as healer that forecloses any possibility of offering care that respects dignity,
21 another in equally good faith understands supporting a patient's request for aid in hastening a
22 foreseen death to be an expression of care and compassion.

23
24 RECOMMENDATION

25
26 The Council on Ethical and Judicial Affairs has reviewed the literature and received thoughtful
27 input from numerous individuals and organizations to inform its deliberations, and is deeply
28 grateful to all who shared their insights. CEJA engaged in extensive, often passionate discussion
29 about how to interpret the *Code of Medical Ethics* in light of ongoing debate and the irreducible
30 differences in moral perspectives identified above. The council recognized that supporters and
31 opponents share a fundamental commitment to values of care, compassion, respect, and dignity, but
32 diverge in drawing different moral conclusions from those underlying values in equally good faith.
33 The council further recognized that medicine must learn from experience of physician-assisted
34 suicide, and must ensure that, where the practice is legal, safeguards are improved.

35
36 After careful consideration, CEJA concludes that in existing opinions on physician-assisted suicide
37 and the exercise of conscience, the *Code* offers guidance to support physicians and the patients
38 they serve in making well-considered, mutually respectful decisions about legally available options
39 for care at the end of life in the intimacy of a patient-physician relationship.

40
41 The Council on Ethical and Judicial Affairs therefore recommends that the *Code of Medical Ethics*
42 not be amended, that Resolutions 15-A-16 and 14-A-17 not be adopted and that the remainder of
43 the report be filed.

Fiscal Note: None.

REFERENCES

1. Council on Ethical and Judicial Affairs. Decisions near the end of life. *JAMA* 1992;267:2229–2233.
2. Vamos MJ. Physician-assisted suicide: saying what we mean and meaning what we say. *ANZJP* 2012;46:84–86.
3. Herx L. Physician-assisted death is not palliative care. *Curr Oncol* 2015;22:82–83.
4. American Association of Suicidology. "Suicide" Is Not the Same as "Physician Aid in Dying". November 12, 2017. Available at <http://www.suicidology.org/Portals/14/docs/Press%20Release/AAS%20PAD%20Statement%20Approved%2010.30.17%20ed%2010-30-17.pdf>. Accessed July 10, 2018.
5. Quill TE, Back AL, Block SD. Responding to patients requesting physician-assisted death: physician involvement at the very end of life. *JAMA* 2016;315:245–246.
6. American Academy of Hospice and Palliative Medicine. *Statement on Physician-Assisted Dying*. June 24, 2016. Available at <http://aahpm.org/positions/pad>. Accessed February 3, 2017.
7. Quill TE, Cassel CK. Professional organizations' position statements on physician-assisted suicide: a case for studied neutrality. *Ann Intern Med* 2003;138:208–211.
8. Sulmasy DP, Finlay I, Fitzgerald F, et al. Physician-assisted suicide: why neutrality by organized medicine is neither neutral nor appropriate. *J Gen Intern Med*. 2018;33(8):1394–1399.
9. Johnstone M-J. Organization position statements and the stance of "studied neutrality" on euthanasia in palliative care. *J Pain Symptom Manage* 2012;44:896–907.
10. Crigger BJ, McCormick PW, Brotherton SL, Blake V. Report by the American Medical Association's Council on Ethical and Judicial Affairs on physicians' exercise of conscience. *J Clin Ethics* 2016;27:291–226.
11. Chochinov HM. Physician-assisted death in Canada. *JAMA* 2016;315:253–254.
12. Battin MP, van der Heide A, Ganzini L, van der Wal G, Onwuteaka-Philipsen B. Legal physician-assisted dying in Oregon and the Netherlands: evidence concerning the impact on patients in vulnerable groups. *J Med Ethics* 2007;33:591–597.
13. Finlay IG, George R. Legal physician-assisted suicide in Oregon and The Netherlands: evidence concerning the impact on patients in vulnerable groups—another perspective on Oregon's data. *J Med Ethics* 2010;37:171–174.
14. Golden M, Zoanni T. Killing us softly: the dangers of legalizing assisted suicide. *Disability and Health Journal* 2010;3:16–30.
15. U.S. Conference of Catholic Bishops. *Assisted Suicide Laws in Oregon and Washington: What Safeguards?* September 22, 2016. Available at http://www.usccb.org/search.cfm?site=newusccb&proxystylesheet=newusccb_frontend&q=assisted+suicide&lang=eng. Accessed October 27, 2016.
16. Ganzini L, Back AL. The challenge of new legislation on physician-assisted death. *JAMA Intern Med* 2016;176:427–428.
17. Lerner BH, Caplan AL. Euthanasia in Belgium and the Netherlands: on a slippery slope? *JAMA Intern Med* 2015;175:1640–1641.
18. Aviv R. The death treatment. *New Yorker*;2015:June 22.
19. Miller DG, Kim SYH. Euthanasia and physician-assisted suicide not meeting due care criteria in the Netherlands: a qualitative review of review committee judgments. *BMJ Open* 2017;7:e017628.
20. Doernberg SN, Peteet JR, KIM SYH. Capacity evaluation of psychiatric patients requesting assisted death in the Netherlands. *Psychosomatics* 2016;57:556–565.
21. Gostin LO, Roberts AE. Physician-assisted dying: a turning point? *JAMA* 2016;315:249–250.

22. Nguyen HQ, Gelman EJ, Bush TA, Lee JA, Kanter MH. Characterizing Kaiser Permanente Southern California's experience with the California End of Life Option Act in the first year of implementation [research letter]. *JAMA Intern Med* 2017;December 26.
23. Li M, Watt S, Escaf M, et al. Medical assistance in dying: implementing a hospital-based program in Canada. *N Engl J Med* 2017;376:2082–2088.
24. Institute of Medicine. *Dying in America: Improving Quality and Honoring Individual Preferences near the End of Life*. Washington, DC: The National Academies Press; 2015.
25. National Academies of Science, Engineering and Medicine. *Physician-Assisted Death: Scanning the Landscape and Potential Approaches*. Available at <https://www.nap.edu/download/25131#>. Accessed August 20, 2018.
26. Dzung E. Can growing popular support for physician-assisted death motivate organized medicine to improve end-of-life care? *J Gen Intern Med*. 2018;33:1209–1211.
27. Quill TE. Doctor, I want to die. will you help me? *JAMA* 1993;270:870–873.
28. Petrillo LA, Dzung E, Smith AK. California's End of Life Option Act: opportunities and challenges ahead. *J Gen Intern Med* 2016;31:828–829.